

Label image proudly supplied by

Infopest
 infopest@dpi.qld.gov.au



500 mL

For the treatment and control of ivermectin sensitive internal and external parasites of cattle and pigs

Active Constituent: 10 mg/mL IVERMECTIN

ANTIPARASITIC INJECTION FOR CATTLE AND PIGS

BOMEECTIN

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS

BEFORE OPENING OR USING

FOR ANIMAL TREATMENT ONLY

BOMEECTIN

ANTIPARASITIC INJECTION FOR CATTLE AND PIGS

Active Constituent: 10 mg/mL IVERMECTIN

For the treatment and control of ivermectin sensitive internal and external parasites of cattle and pigs

BOMAC

500 mL

BOMEECTIN
 ANTIPARASITIC INJECTION FOR CATTLE AND PIGS

BOMAC 500 mL

INDICATIONS
 BOMEECTIN when used at the recommended dose level of 1 mL per 50 kg bodyweight (cattle) or 1 mL per 33 kg bodyweight (pigs) provides control of the following parasites.

CATTLE:
 Gastrointestinal roundworms: adult and immature - *Ostertagia ostertagi* (including third and fourth stage larvae and inhibited immatures), *O. lyrata*, *Haemonchus placei* (including third and fourth stage larvae), *Trichostrongylus axei*, *T. colubriformis*, *Cooperia* spp. (including third and fourth stage larvae), *Bunostomum phlebotomum* (including third and fourth stage larvae) and adults of *Nematodirus spaltzeri*, *Nematodirus helveticus*, *Strongyloides papillosus*, *Toxocara vitulorum* and *Trichuris* spp.
 Lungworms: adult and immature - *Dictyocaulus viviparus* (including inhibited stages).

Eyeworms: adult *Thelazia* spp.
 Sucking Lice: *Linognathus vituli*, *Haematopinus eurysternus* and *Solenopotes capillatus*. Does not provide full control of the cattle biting louse *Bovicola (Damastis) bovis*.
 Sorew Worm Fly: parasitic larval stages of *Chrysomya bezziana* (not reported in Australia but present in Papua New Guinea).
 Mites: *Sarcoptes scabiei* var. *bovis*.
 Cattle Tick: *Boophilus microplus*.

Persistent Activity
 At the recommended dose rate, BOMEECTIN effectively controls infections with *Ostertagia* spp. and *Cooperia* spp. acquired up to 7 days after treatment, and *Dictyocaulus viviparus* and *Chrysomya bezziana* acquired up to 14 days after treatment. When used as directed for control of internal parasites of cattle, BOMEECTIN concurrently aids in control of:

Mites: *Chorioptes bovis*.

PIGS:
 Gastrointestinal roundworms: *Ascaris suum* (adults and L4 immature stage), *Hyostrogylus rubidus* (adults and L4 immature stage), *Oesophagostomum* spp. (adults and L4 immature stage), *Strongyloides ransomi* (adults), *Trichuris suis* (adults).
 Kidney worms: *Stephanurus dentatus* (adults and L4 immature stage).
 Lungworms: *Metsynstrongylus* spp. (adults).
 Lice: *Haematopinus suis*.
 Mites: *Sarcoptes scabiei* var. *suis*.

BOMEECTIN given to the sow 7 to 14 days before farrowing effectively controls transmission via the milk of *Strongyloides ransomi* infections to piglets.

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.

DIRECTIONS FOR USE

Restraints
 Do not use in lactating cows within 28 days of calving where milk or milk products may be used for human consumption.

CATTLE:
 Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers etc). Do not underdose. A representative sample of cattle should be weighed before treatment either with scales or a weighband. The recommended dose is 1 mL BOMEECTIN per 50 kg liveweight (equivalent to 200 µg ivermectin per kg liveweight) by subcutaneous injection. Do not use intravenously or intramuscularly.

Liveweight (kg)	Dose Volume (mL)	Cattle treated per 500 mL pack
Up to 100	2	250
101 - 150	3	166
151 - 200	4	125
201 - 250	5	100
251 - 300	6	83
301 - 350	7	71
351 - 400	8	62
401 - 450	9	55
451 - 500	10	50
501 - 550	11	45
551 - 600	12	41
601 - 650	13	38

Heavier animals (eg mature bulls) should receive an additional 1 mL for each 50 kg over 650.

PIGS:
 Dose animals according to the heaviest animal by liveweight in each group (sows, gilts, boars etc). Do not underdose. Pigs should be weighed. In young pigs, especially those below 16 kg, dosing accurately is important. The recommended dose is 1 mL per 33 kg liveweight (equivalent to 300 µg ivermectin per kg liveweight) by subcutaneous injection in the neck. Do not use intravenously or intramuscularly.

Liveweight (kg)	Dose Volume (mL)	Pigs treated per 500 mL pack
16	0.5	1000
33	1	500
50	2	250
100	3	166
133	4	125
166	5	100
200	6	83

Heavier animals (eg mature boars) should receive an additional 1 mL for each 33 kg over 200 kg.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:
 MEAT: DO NOT USE in pigs or cattle less than 28 days before slaughter for human consumption.
 MILK: When milk or milk products are to be used for human consumption DO NOT USE in cows within 28 days prior to calving or during lactation.

EXPORT SLAUGHTER INTERVAL (ESI) (CATTLE): DO NOT SLAUGHTER cattle for export within 42 days of treatment.

SAFETY DIRECTIONS:
 Poisonous if swallowed. Avoid contact with skin and clothing.
FIRST AID
 If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131126. For further information, refer to the Material Safety Data Sheet.

ENVIRONMENTAL SAFETY:
 Studies indicate that when ivermectin comes in contact with the soil it readily and tightly binds to the soil and becomes inactive over time. Drug containers and any residual contents should be disposed of safely in an approved landfill as free ivermectin may adversely affect fish and certain water-borne organisms.

DO NOT CONTAMINATE DAMS, RIVERS OR STREAMS.

Distributed by:
 Bomac Animal Health Pty Limited (ABN 77 084 246 206)
 8 Apollo Avenue West Pymble NSW 2073
 Orders & Product Information Call 02 9987 4922
 Website: www.bomac.net.au

DISPOSAL:
 Dispose of empty containers and outer packaging by wrapping with paper and putting in garbage.

STORAGE:
 Store below 25°C (Air Conditioning).
 Store bottle in carton to protect from light.

APVMA Approval No. 56848/500mL/0606

Batch: _____
 DOM: _____
 Expiry: _____



CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

BOMECTIN

ANTIPARASITIC INJECTION FOR CATTLE AND PIGS

Active Constituent: 10 mg/mL IVERMECTIN
For the treatment and control of ivermectin sensitive
internal and external parasites of cattle and pigs.

**READ THE ENCLOSED
LEAFLET BEFORE USING
THIS PRODUCT.**

Directions for use

Restraints:

DO NOT USE in lactating cows or
within 28 days of calving where
milk or milk products may be
used for human consumption.

Dose: 1 mL per 50 kg bodyweight
(cattle) or 1 mL per 33 kg bodyweight
(pigs).

WITHHOLDING PERIODS

MEAT: DO NOT USE in pigs or cattle
less than 28 days before slaughter
for human consumption.

MILK: DO NOT USE in lactating cows
or within 28 days of calving where
milk or milk products may be used for
human consumption.

**EXPORT SLAUGHTER INTERVAL
(ESI) (CATTLE): DO NOT
SLAUGHTER** cattle for export within
42 days of treatment.

APVMA Approval No:
56846/500mL/0606

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid
contact with eyes and skin and
clothing.

FIRST AID

If poisoning occurs contact a doctor
or Poisons Information Centre.
Phone 131126.

For further information, refer to the
Material Safety Data Sheet.

DISPOSAL

Dispose of empty containers and outer
packaging by wrapping with paper and
putting in garbage.

STORAGE

Store below 25°C (Air Conditioning).
Store bottle in carton to protect from
light. After vial is first broached, use
within 28 days. After that time discard
any unused portion.

Distributed by:

Bomac Animal Health Pty Ltd
ABN 77 084 248 206
8 Apollo Avenue West Pymble NSW 2073
Orders & Product Information
Call 02 9987 4922

Website: www.bomac.net.au

Batch Number:

Expiry Date:

BOMAC

500 mL