



**SUSTAINED ACTION ORAL ANTHELMINTIC AND FLUKICIDE
FOR SHEEP AND LAMBS:**

Contains Closantel (37.5 g/L), a member of the salicylanilide and substituted phenol family of drenches. It is effective against internal parasites sensitive to this family including the following parasites:

- Haemonchus contortus* Barbers pole worm
- Fasciola hepatica* Liver Fluke
- Oestrus ovis* Nasal Bots

WSD CLOSANTEL provides sustained action against the reinfestation of both mature and immature stages of Barbers Pole Worm in sheep for four weeks after dosing.

WSD CLOSANTEL also controls both mature and immature stages of Liver Fluke as well as all stages of Nasal Bots.

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT USE in female sheep which are producing or may in the future produce milk or milk products for human consumption.

ADMINISTRATION:

Shake container well before using this product.

WSD CLOSANTEL can be administered orally through standard accurate drenching equipment. Clean equipment thoroughly before and after use.

DOSE RATES:

Dose sheep orally to the heaviest animal by liveweight in each group (ewes, rams and lambs) and these should be weighed with scales or weighband. Do not underdose.

Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing.

Internal parasites may develop resistance to drenches. Ask your local veterinary adviser for recommended parasite management practices for your area to reduce development of resistance.

It is advisable that a drench resistance test be conducted before any drench is used. If drench failure with this preparation is suspected, seek veterinary confirmation and notify the manufacturer.

Use 1 mL of Closantel/5 kg bodyweight.				
Bodyweight	Dose Volume	20 L Treats	5 L Treats	1 L Treats
Under 15 kg	3.0 mL	6666	1666	333
16-25 kg	5.0 mL	4000	1000	200
26-35 kg	7.0 mL	2857	714	142
36-40 kg	8.0 mL	2500	625	125
41-45 kg	9.0 mL	2222	555	111
46-50 kg	10.0 mL	2000	500	100
51-55 kg	11.0 mL	1818	454	90
56-60 kg	12.0 mL	1666	416	83
61-65 kg	13.0 mL	1538	384	76
66-70 kg	14.0 mL	1428	357	71
71-75 kg	15.0 mL	1333	333	66
Over 75 kg 1 mL per 5 kg of live bodyweight				

WITHHOLDING PERIOD:

MEAT: DO NOT USE less than twenty eight (28) days before slaughter for human consumption.

MILK: DO NOT USE in female sheep which are producing or may in the future produce milk or milk products for human consumption.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 60 days before slaughter for export. The ESI on this label was current at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on Ph: (08) 9277 5013 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

SAFETY DIRECTIONS:

May irritate the eyes and skin. Avoid contact with the eyes and skin. When opening the container and using the product, wear cotton overalls, buttoned to the neck and wrist and elbow length PVC gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre (Ph. 13 11 26).

MATERIAL SAFETY DATA SHEET (MSDS):

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT:

Do not contaminate dams, streams, rivers or other waterways with the chemical or used containers.

DISPOSAL:

Do not dispose of undiluted chemicals on-site. Break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

CONDITIONS OF SALE:

All Western Stock Distributors products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.

WESTERN STOCK DISTRIBUTORS

A.B.N. 58 009 372 525
Koojan Avenue, South Guildford WA 6055
Telephone: (08) 9277 5013

STORAGE:

Store below 30°C (room temperature) in the closed, original container in a dry, well-ventilated area, and protect from sunlight.

APVMA Approval No: 46165/20L/0609 Batch No:

Expiry: