



Low Volume

LEVAMISOLE

ORAL ANTHELMINTIC FOR SHEEP AND CATTLE contains levamisole hydrochloride (80g/L), a member of the levamisole family of drenches. It is effective for the control of levamisole sensitive gastrointestinal roundworms and lungworms including those that are resistant to benzimidazoles.

It is effective against sensitive strains of the following internal parasites:

Sheep and Lamb

<i>Haemonchus contortus</i>	Barbers pole worm
<i>Trichostrongylus axei</i>	Stomach hair worm
<i>Ostertagia spp*</i>	Small brown stomach worm
<i>Trichostrongylus spp*</i>	Black scour worm
<i>Cooperia spp</i>	Small intestinal worm
<i>Nematodirus spp</i>	Thin necked intestinal worm
<i>Strongyloides papillosus</i>	Intestinal threadworm
<i>Bunostomum spp</i>	Hookworm
<i>Oesophagostomum columbianum</i>	Nodule worm
<i>Oesophagostomum venulosum</i>	Large bowel worm
<i>Chabertia ovina</i>	Large mouthed bowel worm
<i>Dictyocaulus filaria</i>	Large lungworm

Cattle

<i>Haemonchus placei</i>	Barbers pole worm
<i>Trichostrongylus axei</i>	Stomach hair worm
<i>Trichostrongylus spp</i>	Black scour worm
<i>Bunostomum phlebotomum</i>	Hookworm
<i>Oesophagostomum radiatum</i>	Nodule worm
<i>Cooperia spp</i>	Small intestinal worm
<i>Nematodirus spp</i>	Thin necked intestinal worm
<i>Dictyocaulus viviparus</i>	Lungworm

DIRECTIONS FOR USE: Shake container well before using product.

RESTRAINTS: DO NOT USE in female sheep or cattle which are producing or may in the future produce milk or milk products for human consumption.

ADMINISTRATION: WSD LEVAMISOLE can be administered orally through standard accurate drenching equipment. Clean equipment thoroughly before and after use.

Sheep 1mL/10kg live bodyweight

Bodyweight	Dose Volume	20L Treats	10L Treats	5L Treats
Under 10 kg	1.0mL	20000	10000	5000
11-15 kg	1.5mL	13333	6666	3333
16-20 kg	2.0mL	10000	5000	2500
21-25 kg	2.5mL	8000	4000	2000
26-30 kg	3.0mL	6666	3333	1666
31-35 kg	3.5mL	5714	2857	1428
36-40 kg	4.0mL	5000	2500	1250
41-45 kg	4.5mL	4444	2222	1111
46-50 kg	5.0mL	4000	2000	1000
51-55 kg	5.5mL	3636	1818	909
56-60 kg	6.0mL	3333	1666	833
61-65 kg	6.5mL	3076	1538	769
66-70 kg	7.0mL	2857	1428	714
71-75 kg	7.5mL	2666	1333	666
Over 75kg	1mL per 10kg of live bodyweight			

Cattle 5mL/50kg live bodyweight

Bodyweight	Dose Volume	20L Treats	10L Treats	5L Treats
Under 50 kg	5mL	4000	2000	1000
51-100kg	10mL	2000	1000	500
101-150kg	15mL	1333	666	333
151-200kg	20mL	1000	500	250
201-250kg	25mL	800	400	200
251-300kg	30mL	666	333	166
301-350kg	35mL	571	285	142
351-400kg	40mL	500	250	125
401-450kg	45mL	444	222	111
451-500kg	50mL	400	200	100
501-550kg	55mL	363	181	90
551-600kg	60mL	333	166	83
601-650kg	65mL	307	153	76
651-700kg	70mL	285	142	71
Over 700kg	5mL per 50kg live bodyweight with a maximum of 90mL for animals 900kg and over			

DOSE RATES: Dose mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs); (bulls, cows, steers, calves) and these should be weighed with scales or weighband. Do not underdose.

Where there is a large variation in size within the group, dose rate should be based on label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

WARNING: Internal parasites may develop resistance to drenches (anthelmintics). Ask your local veterinary adviser for recommended parasite management practices for your area to reduce development of resistance.

It is advisable that a drench resistance test be conducted before any drench is used. If drench failure with this product is suspected, seek veterinary confirmation and notify the manufacturer.

PRECAUTIONS: Exercise care in handling weak or pregnant animals to avoid unnecessary stress. DO NOT DRENCH DOGS WITH WSD LV LEVAMISOLE drench.

WITHHOLDING PERIODS: MEAT: DO NOT USE less than three (3) days before slaughter for human consumption.

MILK: DO NOT USE in female sheep or cattle which are producing or may in the future produce milk or milk products for human consumption.

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT slaughter for export 7 days after treatment.

SAFETY DIRECTIONS: Harmful if swallowed. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26.

MATERIAL SAFETY DATA SHEET (MSDS): Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

PROTECTION OF FISH, CRUSTACEANS AND THE ENVIRONMENT: Dangerous to fish. Do not contaminate streams, rivers and waterways with the product or used containers.

DISPOSAL: Do not dispose of undiluted chemicals on-site. Break, crush, puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

STORAGE: Store below 30° (room temperature) in the closed, original container, in a dry, well-ventilated area, and protect from sunlight.

CONDITIONS OF SALE: All Western Stock Distributors products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.

WESTERN STOCK DISTRIBUTORS:

A.B.N. 58 009 372 525
Koojan Avenue, South Guildford WA 6055
Telephone: (08) 9321 2888 Facsimile: (08) 9322 4163
www.westernstock.com.au

APVMA Approval No: 39063/1100

Batch No:

Expiry: